

PIC Nutrient Specifications – 2008

Introduction

This publication provides a summary of nutrient recommendations for PIC pigs. Recommendations are based primarily on published and internal research, research from the University of Missouri and Ajinomoto Heartland with commercial PIC products. The nutrient specifications have been validated in commercial environments. The NRC publication (1998) serves as the basis for certain information. Concepts and the basis for recommendations are discussed in greater detail in other Technical Memos.

Contents:

Dietary Energy

Formulating to a SID Amino Acid Pattern

Camborough® Gilt Product Development Specifications

- Gilt breeding targets: 210 days of age and 300 lb and 1 heat no service.
- Nutrient specification table for gilt development.

Camborough® Sow Product Specifications

- Sow Feeding Milestones
- Daily Nutrient Needs and Diet Specifications: Pregnancy
- Energy and Lysine Needs for Pregnancy Weight Gain
- Daily Nutrient Needs and Diet Specifications: Lactation
- Estimating Farm-specific Lactation Lysine Needs
- Wean 365 pounds of pigs/sow/year

Boar Stud Specifications

- Minimum Feeding Levels
- Growth Rates of Boars Fed to PIC Intake Standards
- Diet Specifications for Stud Boars

Nursery Diet Specifications

- Diet Specifications for a 3-4 Phase Nursery Program
- Example Feeding Program and Feed Budget
- Matching Lysine Need to Different Growth Phases

Grow-Finish Specifications

- Diet Specifications for Optimum Lean Deposition: Gilts
- Diet Specifications for Optimum Lean Deposition: Castrates
- Adjusting Lysine-Calorie Specifications to Lean Content
- Diet Specifications for Poor or No Lean Payment Markets
- Matching Lysine Need to Different Growth Phases
- Wean to Finish ADG > 1.75 lb and F/G < 2.45
- Feeder to Finish ADG > 1.95 lb and F/G < 2.65

Vitamin and Trace Mineral Addition

- Commercial
- Multiplication

Other sections:

- Paylean® specifications
- Alternative feed stuffs
- How nutrition affects yield
- Feed form (pellets vs meal)
- Feeder space

References



Dietary Energy

Energy levels are expressed as Metabolizable energy (ME) in this guide because it is the system most commonly used in North and South America. The ME (and DE) system is adequate for describing available energy for simple diets (Corn-Soy) or for diets of young pigs. However, the Net energy (NE) system is a more accurate method of measuring the available energy when a more diverse set of ingredients are used. NE is in routine use by several feed companies for this reason. Other groups have 'modified' ME values for certain ingredients (eg., soybean meal, wheat midds) to reflect NE relationships. This guide specifies energy as NRC ME, based on a simple diet type that is defined for each phase. Nutrition specialists can then adapt the specified Lysine:ME relationships to their own system. Adjustments in ME for hot environmental temperature or for profit related fat use will require adjusting Lysine levels to keep Lysine:ME constant.

Tables 1, 2 and 3 below are tables that demonstrate the response of PIC pigs to added energy. The first table is the overall means and the second and third show the means of gilts and barrows for ADG and F/G.

Table 1. Effects of added fat to a corn - soybean meal diet^a

| | Added fat, % | | | |
|-----------------|--------------|-------|-------|-------|
| | 1.0 | 2.5 | 4.0 | 5.5 |
| NRC ME, kcal/lb | 1471 | 1503 | 1534 | 1565 |
| ADG, lb | 2.11 | 2.11 | 2.14 | 2.16 |
| ADFI, lb | 5.47 | 5.39 | 5.33 | 5.23 |
| F/G | 2.59 | 2.55 | 2.49 | 2.42 |
| HCW, lb | 214.9 | 217.4 | 216.9 | 218.5 |
| FOM lean, % | 53.1 | 53.1 | 53.3 | 53.3 |

^a A total of 4,900 pigs (PIC 337 x Camborough; initial weight 75 lb) with 50 replications per treatment. There were 100 pens each of barrows and gilts with 24 to 25 pigs per pen.

Table 2. Effects of ADG to added fat to a corn - soybean meal diet

| Weight Range, lb | Gender | Added fat, % | | | |
|------------------|---------|--------------|------|------|------|
| | | 1.0 | 2.5 | 4.0 | 5.5 |
| 78 to 120 | Barrows | 2.03 | 2.06 | 2.09 | 2.08 |
| | Gilts | 2.02 | 2.02 | 2.06 | 2.05 |
| 120 to 165 | Barrows | 2.25 | 2.32 | 2.35 | 2.43 |
| | Gilts | 2.02 | 2.05 | 2.07 | 2.11 |
| 165 to 210 | Barrows | 2.46 | 2.48 | 2.50 | 2.44 |
| | Gilts | 2.17 | 2.19 | 2.21 | 2.19 |
| 210 to 250 | Barrows | 2.01 | 2.13 | 2.09 | 2.18 |
| | Gilts | 2.00 | 1.96 | 1.98 | 1.98 |

Table 3. Effects of F/G to added fat to a corn - soybean meal diet

| Weight Range, lb | Gender | Added fat, % | | | |
|------------------|---------|--------------|------|------|------|
| | | 1.0 | 2.5 | 4.0 | 5.5 |
| 78 to 120 | Barrows | 1.99 | 1.94 | 1.91 | 1.90 |
| | Gilts | 2.05 | 2.02 | 1.93 | 1.94 |
| 120 to 165 | Barrows | 2.46 | 2.31 | 2.27 | 2.18 |
| | Gilts | 2.39 | 2.29 | 2.17 | 2.09 |
| 165 to 210 | Barrows | 2.92 | 2.83 | 2.76 | 2.74 |
| | Gilts | 2.81 | 2.74 | 2.75 | 2.69 |
| 210 to 250 | Barrows | 3.29 | 3.04 | 3.00 | 2.77 |
| | Gilts | 2.99 | 3.10 | 2.99 | 2.97 |

The results from this trial demonstrate that barrows respond to added fat better than gilts. In this trial for every 1% added fat barrows improved ADG by 0.8% and F/G by 2.0%. For gilts, every 1% added fat, ADG improved by 0.3% and F/G by 1.2%.

Pig flow can influence the value of added fat in a production system. A producer with a fixed time in finishing or who is short of finishing space can pay a higher price for added fat. Whereas a producer that is on a variable time system or has adequate finisher space can hold the pigs for 5 to 7 days longer and the value of additional fat in the diet is not as great.

Formulating to an Ideal Amino Acid Pattern

To obtain optimum performance, all amino acids must meet or exceed the requirement. The NRC (1998) has defined the ideal balance of amino acids for each physiological phase. NRC (1998) served as the basis for PIC recommendations for sows but recent research based modifications have been made for lactation sows and growing pigs based on the collaboration work of PIC, Ajinomoto Heartland and the University of Missouri. They are normally expressed in relation to the level of lysine since it is most likely to be first limiting in the diet. Since ingredient

amino acids differ in their digestibility, standardized ileal digestible (SID) amino acid values are preferred when formulating diets that are more complex than corn-soybean meal diets.

This guide specifies the SID lysine requirement, which is satisfactory for corn-soy formulations. The suggested ratio of dietary amino acids for each phase is as follows.

Table 4. Ideal Pattern for breeding herd based on SID Amino Acid Formulations^{a,b}

| Amino Acid | Gestation ^c | | Lactation ^d | | Boar Stud |
|----------------------|------------------------|------|------------------------|------|-----------|
| | Gilt | Herd | Gilt | Herd | |
| Lysine | 100 | 100 | 100 | 100 | 100 |
| Methionine + Cystine | 70 | 70 | 50 | 50 | 70 |
| Threonine | 76 | 76 | 64 | 64 | 74 |
| Tryptophan | 18 | 18 | 18 | 18 | 20 |
| Valine | 68 | 68 | 75 | 75 | 67 |
| Isoleucine | 58 | 58 | 56 | 56 | 58 |

^a Lysine set equal to 100% and other amino acids expressed as a percent of Lysine.

^b Ratio's are appropriate IF applied to Standardized Ileal Digestible (SID) amino acids (Corn-Soy based) or which is suggested for more complex diets.

^c Assumption: Gilt – 300 lb body weight (bw) at breeding and 75 lbs net maternal gain
Sow – 400 lbs bw at breeding and 20 lbs net maternal gain

^d Assumption: 385 lb bw post-farrowing, 22 lb weight loss & 2200-2500 g/d Litter growth.

Table 5. Ideal Pattern for nursery and finishing pigs based on SID Amino Acid Formulations

| Amino Acid | Nursery | Finish | | | | | Paylean® |
|------------|------------|-------------|--------------|---------------|---------------|---------------|---------------|
| | 8 to 50 lb | 50 to 90 lb | 90 to 130 lb | 130 to 180 lb | 180 to 230 lb | 230 to 280 lb | 230 to 280 lb |
| Lysine | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| M + C | 58 | 55 | 55 | 55 | 55 | 55 | 55 |
| Threonine | 60 | 61 | 62 | 63 | 64 | 66 | 68 |
| Tryptophan | 16 | 16 | 16 | 16 | 16 | 16 | 16 |
| Valine | 65 | 68 | 68 | 68 | 68 | 68 | 68 |
| Isoleucine | 55 | 56 | 56 | 56 | 56 | 56 | 56 |

Camborough® Gilt Product Development Specifications.

Gilt nutrition during development has a significant impact on early and lifetime performance of females.

Gilt development and management begins in the early stages of a gilts life and ends when the gilt completes her first lactation. This topic has been discussed by Boyd and Williams at the 2008 Banff conference.



Recommended Developing Gilt Growth and Body Composition Targets are set out below.

- Birth to first service daily gains of 1.40 to 1.45 pounds/day
- Achieve a body weight of 300 pounds at first service (210 days of age)
- Target P2 back fat of 14-16 mm at 1st service (12 mm minimum)

Nutrient requirements for development are found in Table 6.

- The requirements are designed to meet nutrient demands for adequate protein growth and not designed to maximize ADG (as would be the goal of a commercial feeding program.(see Grow finish specifications, "Nutrient levels for optimum lean deposition" for commercial feeding)
- Gilt development energy recommendations are 5-6 % lower than typical commercial diets after the nursery phase.

Vitamin and trace mineral requirements are higher than commercial recommendations in order to prepare the gilt for reproductive function (See Vitamin and trace mineral for Genetic nucleus and multiplication table)

For further information relating the Nutrition management of the developing gilt please refer to the Camborough user guide and product manual published by PIC.

Table 6. PIC Nutrient Specifications for Gilt Development^a

| Nutrient | Unit | Weigh range, lb | | | | |
|-------------------------------|----------------|-----------------|---------|----------|-----------|-----------|
| | | 25 - 50 | 50 - 90 | 90 - 150 | 150 - 210 | 210 - 300 |
| NRC ME | Mcal/lb | 1510 | 1490 | 1490 | 1490 | 1490 |
| SID lysine^b | % | 1.32 | 1.15 | 0.95 | 0.80 | 0.65 |
| Calcium | % | 0.80 | 0.75 | 0.70 | 0.70 | 0.70 |
| aPhosphorus | % | 0.42 | 0.36 | 0.35 | 0.35 | 0.35 |

^a Amounts are expressed as the concentration per pound of complete diet.

^b Lysine levels are provided as the bases for normal lean:fat ratios (SID = standard ileal digestible and a = available). Other nutrients are the nutrients most related to proper bone development. See Vitamin and Trace mineral addition rates for Multiplication.

Note: Goal for first service is 210 days, 300 lb and one heat no service recorded. All targets need to be accomplished for proper gilt development (See 2007 PIC gilt and sow manual). Depending on the environment, square footage, etc, nutrition (especially energy and amino acids) will need to be adjusted.

Adjustments will vary from producer to producer depending on gilt growth.

Camborough® Sow Product Specifications

The sow feeding program is a phase feeding system. Feeding involves a gestation diet and 1 or preferably 2 lactation diets. Pregnant sow feeding requires (1) feeding the proper amount of diet to meet energy needs and, (2) then matching nutrient level to the amount fed so that daily nutrient need is met (in g/day). The minimum amount of energy and nutrients is provided in Table 8. These amounts are driven by the amount of growth that we want to achieve as shown in Table 9. Lactation is especially demanding and requires full-feeding beginning day 4. The recommended feeding program of lactating sows is to feed 4 lb on day 1 and 6 lb on day 2 and 6 lb on day 3. On day 4 the sows need to be on full feed by always keeping fresh feed in front of the sow at all times. PIC research has proved that prolific PIC gilts and sows produce more milk which requires improved diets (Boyd et al., 2000). Further, the typical lactation diet is inadequate for gilts so a second diet is advised when feasible especially for start up farms with all gilts and parity segregation farms (Table 3). The main difference between gilt or P1 lactation diet and older sows is the amount of digestible lysine. There is also a cost savings involved with parity segregation feeding with approximate savings of \$3.40 per sow per year.




Table 7. Sow Feeding Milestones (Concepts applied from T. Update on Feeding Prolific Sows).

| Milestone | Amount, lbs ^a | Feed Type |
|-------------------------------------|-----------------------------|------------|
| Pre-breeding Gilt | Full-feed (pre-breed flush) | Gestation |
| Gestation to 5 d Post-breed | | |
| Gilts | 4.5 | Gestation |
| Sows | 4.5 | Gestation |
| Very Thin Sows | 6.5 | Gestation |
| Gestation: 5 d to 90 ^b | per body condition score | Gestation |
| Gestation: 90 d to 116 ^c | 6.0 – 6.5 | Gestation |
| Pre-farrow 2-4 d | 4.0 – 5.0 | Lactation |
| Lactation: 1 d Post - farrow | 4 | Lactation |
| Lactation: 2 and 3 d Post - farrow | 6, 6 | Lactation |
| Lactation: 4 d to weaning | Full-feed | Lactation |
| Weaning through breeding | Full-feed | Gestation* |

^a Assumes 1465 Kcal NRC ME/lb for Gestation and 1500 Kcal NRC ME/lb for Lactation diets.

^b Minimum feed levels to reclaim body fat reserves are shown in Table 4. Objective is to reclaim body reserves (fat, protein, bone minerals) by 60 d pregnant.

^c Average gestation length, 116 d.

- Where possible the weaned sow should be fed 3 to 4 times per day; the pregnant sow should be fed twice daily, where feasible.

Daily Nutrient Needs and Diet Specifications for Gestation

Nutrient specifications for pregnancy are presented in Table 8. Energy and Lysine needs are based on suggested levels of growth in Table 9. Gilt specifications can be used when stocking units. Specifications for herds are a compromise between needs for gilts vs. older sows. Gilts are challenged to a greater extent during lactation than sows. Feed gilts 4.5-5.0 lbs/day until day 90 of pregnancy, this will allow for significant growth to prepare for the rigor of first lactation.

Table 8. Recommendations for Gestating PIC Gilts and Sows

| Nutrient | Unit | Gilt | Herd |
|--------------------------|-------------|-------------|-------------|
| Net wt gain ^a | lb | 75 | Table 2 |
| P2 fat endpoint | mm | 14-16 | 14-16 |
| NRC ME ^b | Kcal/lb | 1465 | 1465 |
| NRC ME ^b | Kcal/d | 6886 | 7179 |
| Calcium | grams/day | 19.4 | 20.0 |
| aPhosphorus ^c | grams/day | 8.8 | 8.9 |
| Crude Fiber | % | 4 to 6 | 4 to 6 |
| Calcium | % | 0.95 | 0.90 |
| aPhosphorus ^c | % | 0.43 | 0.40 |
| Added salt | % | 0.45 | 0.45 |
| Max L – lysine | % | 0.15 | 0.15 |
| Feed intake ^d | lb | > 4 | > 4 |
| SID lysine ^c | % | 0.62 | 0.62 |
| SID lysine | | | |
| d 0 to 90 | grams/day | 14 | 14 |
| d 90 to 116 | grams/day | 18 | 18 |
| d 90 to farrow | | | |
| NRC ME ^b | Kcal/d | 9156 | 9156 |

^a Net tissue growth by the sow declines with each parity. See Table 9 for suggested gain and feed level.

^b NRC ME is based on a Corn-Soy diet with 15% wheat midds.

^c SID = standard ileal digestible. Formulate to SID lysine levels.

The 0.62% SID lysine level is based on 4.5 lb per day intake.

^d Do not feed less than 4 lb per day at any given time period. This will go below maintenance requirement.

Energy and Lysine Needs for Pregnancy Weight Gain^a

Daily energy and lysine levels in Table 8 are minimums and assume proper body condition. Estimates are driven by the amount needed for suggested growth by parity. Larger females require more energy for maintenance, but this increase is off-set by declining maternal gain. This is for gilts and sows from breeding to day 90 of gestation only after 90 days an increase of feed by 1 to 2 lbs is recommended.

Table 9. Energy and lysine needs for gestation weight gain

| Body Weight at Mating, lbs. | LITTER NO. | Suggested Net Gain, lbs | Feed Intake, lbs/d | NRC ME Kcal/d | SID Lysine g/d |
|-----------------------------|-----------------|-------------------------|--------------------|---------------|----------------|
| 280 | 1 st | 75 | 4.7 | 6886 | 12.3 |
| 330 | 2 nd | 75 | 5.0 | 7325 | 12.5 |
| 385 | 3 rd | 55 | 4.9 | 7275 | 12.3 |
| 420 | 4 th | 45 | 4.9 | 7205 | 11.9 |
| 440 | 5 th | 35 | 4.7 | 7070 | 11.5 |
| 450 | 6 th | 30 | 4.8 | 6930 | 11.4 |
| 460 | 7 th | 20 | 4.6 | 7035 | 11.1 |

^a Computed using the 1998 NRC Gestation Model. Assumed 1465 Kcal NRC ME/lb diet and 12 fetuses and also assumes sows needed 2 mm of back fat. SID lysine estimated by 86% of total lysine.

Improve satiety in gestation sows with fiber

Since feed intake is restricted during gestation, to prevent excess body weight gain, sows can often appear restless. For this reason, ingredients that have high levels of insoluble non-starch components are often used (hulls from soybeans and oats; alfalfa) at the rate of 15 to 20% of the diet. This tends to increase satiety and shifts some of the digestion to the cecum from which volatile fatty acids arise.

Below is an example of a gestation diet formulated to a SID lysine 0.62%. The use of fiber is recommended and in this example DDGS and wheat midds are used, other fibrous ingredients that will work is soy hulls, wheat shorts, and oat hulls.

Table 10. PIC Gestation Example Diet

| Ingredient | Pounds |
|-------------------------|--------|
| Corn | 1160.0 |
| SBM | 170.0 |
| DDGS | 300.0 |
| Wheat Midds | 295.0 |
| Dicalcium Phosphate | 19.0 |
| Limestone | 33.0 |
| Salt | 9.0 |
| L-lysine | 3.0 |
| L-threonine | 0.5 |
| VTM + Choline + Phytase | 10.5 |

Daily Nutrient Needs and Diet Specifications for Lactation

Nutrient specifications are presented in Table 11. Daily ME intake is the multiple of (1) expected daily feed intake (based on intakes achieved in commercial sow systems) and (2) the ME content of a Corn-Soy diet. The level of Lysine for gilts is based on PIC research to optimize second litter-size (20 d lactation). Two lactation diets are recommended when stocking units and should be a consideration when designing new units. A single herd diet will increase loss of body protein in gilts, which may result in a second litter-size 'dip or plateau'.

The goal for lactation is to wean 365 pounds of pigs/sow/year. By following the nutrition specs and working with the PIC Technical Service group, this will be achieved. For more information on achieving these and other reproductive targets, please refer to the 2007 PIC gilt and sow technical manual.

13.5 lbs x 11 pigs weaned x 2.46 litters/sow/year = 365 lbs

Table 11. Recommendations for Lactating PIC Gilts and Sows^a

| Nutrient | Unit | Gilt | Sow | Herd |
|------------------------------|-----------|-------|-------|-------|
| Net weight body loss | % | < 10 | < 10 | < 10 |
| Fat Loss, max. | mm | 0 - 2 | 0 - 2 | 0 - 2 |
| Litter growth | lb/day | 4.95 | 5.28 | 5.12 |
| | grams/day | 2250 | 2400 | 2325 |
| NRC ME | Kcal/lb | 1550 | 1510 | 1525 |
| Ave feed intake ^b | lb/day | 11.0 | 12.5 | 12.0 |
| | kg/day | 5.0 | 5.7 | 5.5 |
| NRC ME | Kcal/day | 17050 | 18875 | 18300 |
| Calcium | grams/day | 47 | 51 | 47 |
| aPhosphorus | grams/day | 21 | 23 | 21 |
| SID lysine | grams/day | 63 | 49 | 55 |
| SID lysine | % | 1.22 | 0.87 | 0.95 |
| Calcium | % | 0.95 | 0.90 | 0.95 |
| aPhosphorus | % | 0.43 | 0.40 | 0.43 |
| Added salt | % | 0.45 | 0.45 | 0.45 |
| Max L – lysine | % | 0.30 | 0.30 | 0.30 |

^a About 85-90% energy balance with litter growth rates shown and 20 d lactation (1998 NRC model).

^b Average feed intake over suckling period not maximum intake achieved within the period.

How does the SID lysine requirement differ with each parity?

Srichana et al., 2006 demonstrated with PIC Camborough sows the SID lysine in lactation decreases as the sow matures (see figure 1 below).


P1: 1.20% SID lysine

P2: 0.92% SID lysine

P3: 0.82% SID lysine

P4: 0.72% SID lysine

SID Lysine Requirement of PIC Lactating Sows



Below is an example of a diet for PIC sows formulated to a SID lysine level of 0.95%

Table 12. PIC Lactation Example Diet

| Ingredient | Pounds |
|-------------------------|--------|
| Corn | 1337.4 |
| SBM | 464.0 |
| DDGS | 100.0 |
| CWG | 20.0 |
| Dicalcium Phosphate | 28.4 |
| Limestone | 26.4 |
| Salt | 9.0 |
| L-lysine | 4.0 |
| L-threonine | 0.8 |
| VTM + Choline + Phytase | 10.0 |

Estimating Farm-specific Lactation Lysine Needs

The farm-specific lysine level for lactation depends on the actual litter growth rate and average lactation feed intake by sows. The daily lysine requirement is driven strictly by rate of litter growth and this can vary with health and thermal stress. This needs to be matched with the level of feed consumed. Table 11 could be used to derive farm-specific lysine needs. It is appropriate for sows but not for gilts since a higher lysine level is needed for gilts to maximize second litter size than is required to support milk production (see Current Concepts for Feeding Prolific Sows).

Where possible the provision of a gilt lactation diet should be used; where this is not possible top-dressing with soybean meal is recommended.

Table 13. Lactation Lysine Level Varies with Litter Growth Rate and Sow Feed Intake (for sow only)

| Litter ^a Growth, lbs/day | Litter ^b Wean Wt, lbs | Average Feed Intake, lbs/day ^d (SID lysine, %) | | | | | | SID Lysine ^{c,d} g/day |
|---|--|--|------|------|------|------|------|---------------------------------------|
| | | 9 | 10 | 11 | 12 | 13 | 14 | |
| 3.65 | 110 | 0.86 | 0.77 | 0.69 | 0.65 | 0.60 | -- | 34 |
| 4.15 | 120 | 0.95 | 0.86 | 0.77 | 0.73 | 0.69 | 0.60 | 40 |
| 4.60 | 130 | 1.08 | 0.95 | 0.86 | 0.82 | 0.73 | 0.69 | 44 |
| 5.10 | 140 | -- | 1.08 | 0.99 | 0.90 | 0.82 | 0.77 | 49 |
| 5.55 | 150 | -- | -- | 1.08 | 0.99 | 0.90 | 0.86 | 54 |
| 6.00 | 160 | -- | -- | 1.17 | 1.08 | 0.99 | 0.92 | 58 |

- ^a Assumes 11 pigs with an average birth weight of 3.3 lb/pig. LGR is calculated as follows: ((Litter-Wean Weight – (No. nursed x 3.3 lbs/pig)) ÷ Lactation length, days.
- ^b Litter wean weight applicable to a 21d lactation. Adjustment factors for different wean ages were suggested by NSIF, 1987.
- ^c Calculated from updated Pettigrew equation (Boyd, et al., 2000a) and assumes that Lysine need is not strictly related to ME intake. The equation is based on a linear relationship between LGR and Lysine need (g/d) to support milk production.
- ^d Feed Intake assumes 1500 Kcal NRC ME/lb. SID lysine is based on 86% of total lysine.

Boar Stud Specifications

Energy needs to support target weight gains (Table 14/ Figure 2) without compromising sperm output have been calculated and validated in AI studs (Technical memo 142.) Higher levels may result in maximum sperm output. Very little information exists on which to base Nutrient specifications. Those in Table 15 are used by PIC and given for reference only. Energy and amino acid levels are based on limited University research.




Table 14. Minimum Feed Level in Relation to Body Weight^a

| Body Weight Lbs | Body Weight kg | Total Kcal ME/day ^b | Feed lbs/day | Feed kg/day |
|--------------------|-------------------|-----------------------------------|-----------------|----------------|
| < 350 | < 159 | 7200 | 5.0 | 2.3 |
| 350 | 159 | 7920 | 5.5 | 2.5 |
| 450 | 205 | 8640 | 6.0 | 2.7 |
| 550 | 250 | 9505 | 6.6 | 3.0 |
| 650 | 295 | 10370 | 7.2 | 3.3 |
| 750 | 341 | 11230 | 7.8 | 3.5 |

^a Adapted from PIC Technical Memo 142. Assumes ambient temperature of 62-65°F.

^b Based on a dietary energy density of 1440 Kcal ME/lb.



^aAdapted from PIC Technical Memo 142. 1 kg of Gain = 2.2 lbs.

Table 15. Boar Minimum Diet Specifications^a

| Nutrient | Unit | |
|---------------------------|-------------|------------|
| NRC ME | Kcal/lb | 1400 |
| Protein | % | 16 |
| Fiber | % | 4.5 to 6.0 |
| SID lysine ^b | % | 0.62 |
| Calcium | % | 0.85 |
| T Phosphorus ^b | % | 0.75 |
| aPhosphorus ^b | % | 0.40 |
| Added salt | % | 0.45 |
| Linoleic acid | % | 1.90 |

^a Amount / lb of complete diet assuming feed level in Table 1.

See Vitamin & Trace mineral section for specifications.

^b T = total, a = available, SID = Standardized ileal digestible.
Formulate to either Total or SID levels but not both.

Diet specifications are based on the minimum feed levels shown in Table 14 above. Vitamin and trace mineral levels are presented in the final section for all phases of production including the boar stud. Organic mineral sources are recommended especially for selenium, copper, zinc and manganese.

Nursery Diet Specifications

Nutrient specifications are shown in Table 16. A feeding program and budget is illustrated in Table 17. They are based on extensive research information from production companies and published literature. A soybean maximum is suggested for each phase as a starting point for commercial conditions. Higher levels may result in 'looseness' but should be tested so that a farm-specific maximum can be set.

Diets must be matched to body weight and the correct amount fed within each phase. Feeding 2 diets simultaneously within a weaned group is advised through phase 2 for optimum growth and livability. Typical weight variation at weaning means that feeding to the average pig weight will result in feeding the diet that matches requirements for only 65-70% of the pigs (Table 17, footnote b). This will result in increased variation in nursery end weight.

Table 16. Diet Specifications for a 3 to 4 Phase Nursery Program.

| Nutrient | Unit | Early Wean 8-12 lbs | Phase 2 12-16 lbs | Phase 3 16-25 lbs | Phase 4 25-50 lbs |
|------------------------------------|---------|------------------------|----------------------|----------------------|----------------------|
| Growth rate | lb/d | 0.34 | 0.50 | 0.90 | 1.45 |
| Feed Intake ^a | lb/d | 0.35 | 0.58 | 1.18 | 2.20 |
| Feed:Gain | Ratio | 1.03 | 1.16 | 1.31 | 1.52 |
| NRC ME | Kcal/lb | 1550 | 1550 | 1530 | 1525 |
| Lactose ^b | % | 20.0 | 15.0 | 7.5 | 0.0 |
| Total Fat | % | 3 to 6 | 3 to 6 | 2 to 4 | 2 to 4 |
| SID lysine ^c | % | 1.46 | 1.46 | 1.42 | 1.30 |
| SID Meth + Cystine:lysine | Ratio | 0.58 | 0.58 | 0.58 | 0.58 |
| SID Threonine:lysine | Ratio | 0.60 | 0.60 | 0.60 | 0.60 |
| SID Tryptophan:lysine | Ratio | 0.16 | 0.16 | 0.16 | 0.16 |
| SID Valine:lysine | Ratio | 0.65 | 0.65 | 0.65 | 0.65 |
| SID Isoleucine:lysine ^e | Ratio | 0.55 | 0.55 | 0.55 | 0.55 |
| Calcium | % | 0.85-0.95 | 0.85-0.95 | 0.80-0.90 | 0.80-0.90 |
| aPhosphorous | % | 0.55 | 0.52 | 0.40 | 0.40 |
| aPhos:ME | g/Mcal | 1.58 | 1.50 | 1.16 | 1.16 |
| Sodium | % | 0.35-0.85 | 0.35-0.60 | 0.25-0.40 | 0.25-0.40 |
| Chloride | % | 0.40 | 0.40 | 0.36 | 0.36 |
| Added salt | % | 0.20 | 0.25 | 0.40 | 0.40 |
| Potassium | % | 0.70 | 0.70 | 0.65 | 0.65 |
| Soybean meal ^d | % | 15 | 18 | 28 | 28-32 |

^a Average Intake shown for 25-50 lb pig assumes pelleted. Add 5% for grind and mix.

^b Lactose or equivalent sugars. Sucrose can replace 50% Lactose after 5-7 d of feeding to induce digestive enzyme.

^c T = Total and SID = standard ileal digestible.

^d Suggested levels for commercial production and good to high health. High health pigs can tolerate higher levels of SBM (16-25lbs,30% ; 25-50lbs,32%)

^e Diet with < 1% blood cells.

Feed Budgeting. Providing the proper amount of each diet and avoiding excesses is important to economics. Feed budgets allow a producer to predict and control feed cost. The budgeted amount should be verified at the outset by weighing a representative sample of pigs at each diet switch. The actual amount fed of each diet should be within 5% of target.

An example feed budget is shown in Table 17 for three weaning scenarios. The budgeted amount of feed per phase is driven by feed conversion (FCR) data in Table 16. The percentage of pigs that are expected to be smaller than the average is also considered (Table 17, footnote b). A change in FCR will result if diet energy density is different than shown in Table 16 (especially for 25-50 lbs phase) so the budgeted amount must be changed to reflect the difference.

Table 17. Example Feeding Program and Feed Budget^{a,b}

| Ave Wean Age | days | 14 | 17 | 20 |
|------------------------|---------------|----------------------------|-------------|-------------|
| Ave Wean Weight | lb | 10.5 | 12.0 | 13.5 |
| Diet Phase | Weight | Feed Budget per pig | | |
| Phase 1 | 8 - 12 | 2.0 | 1.0 | 1.0 |
| Phase 2 | 12 - 16 | 4.0 | 4.0 | 4.0 |
| Phase 3 | 16 - 25 | 12.0 | 12.0 | 12.0 |
| Phase 4 | 25 - 50 | 40.0 | 40.0 | 40.0 |

| | | | | |
|-------------------|--------------|-------------|-------------|-------------|
| Total Feed | lbs | 58.0 | 57.0 | 56.0 |
| Total Gain | lbs | 39.5 | 38.0 | 36.5 |
| Feed:Gain | Ratio | 1.47 | 1.50 | 1.56 |

^a Budget assumes 50 lb end weight for Nursery and FCR shown in Table 1.

^b Early Wean diet allocated assuming FCR = 1.05 and a Population percent by wean weight as follows:
 Population %, Avg 12 lbs wean weight: $\frac{9 \text{ lbs}}{5\%}$ $\frac{10 \text{ lbs}}{10\%}$ $\frac{11 \text{ lbs}}{20\%}$ $\frac{12 \text{ lbs}}{30\%}$ $\frac{13 \text{ lbs}}{20\%}$ $\frac{14 \text{ lbs}}{10\%}$ $\frac{15 \text{ lbs}}{5\%}$

Grow-Finish Specifications: Nutrient Levels for Optimum Lean Deposition

Nutrient specifications in Tables 18 and 19 are for lean growth optimization. Performance was determined under conditions of high health, thermoneutral environment and with corn-soy diets.

Lysine specifications are presented as grams per Mcal of NRC ME. Formulating diets of growing pigs to grams of Lysine/day can be misleading (discussed in Grow-Finish Technical Update). The dietary energy level used in the recommendations is 1500 or 1600 Kcal NRC ME/lb since it approximates the corn-soy diet. Actual dietary energy levels require a number of considerations that are specific to market and environment (see Usry et al., 1997). The available phosphorus curve (g P/Mcal NRC ME) is higher than the NRC for Nursery and especially for Grow-Finish phases because levels have not proven adequate to withstand the force imposed on bones during stunning in commercial plants (as compared to University Labs).

Table 18. PIC Gilt Specifications^a

| Gilts | | Phase of growth, lb | | | | | | | | | |
|--|---------|---------------------|------|----------|------|-----------|------|-----------|------|------------------------|------|
| Item | Unit | 50 - 90 | | 90 - 130 | | 130 - 180 | | 180 - 230 | | 230 - 280 ^e | |
| Growth rate | lb/d | 1.78 | | 1.91 | | 2.15 | | 2.08 | | 1.98 | |
| Feed intake | lb/d | 3.00 | | 4.28 | | 5.59 | | 5.72 | | 5.98 | |
| Feed:Gain | Ratio | 1.69 | | 2.24 | | 2.60 | | 2.75 | | 3.02 | |
| NRC ME | Kcal/lb | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 |
| Added fat | % | 0 | 5 | 0 | 5 | 0 | 5 | 0 | 5 | 0 | 5 |
| SID lysine/Mcal ME ^b | g/Mcal | 3.17 | 3.17 | 2.76 | 2.76 | 2.39 | 2.39 | 2.11 | 2.11 | 1.96 | 1.96 |
| SID lysine ^c | % | 1.05 | 1.12 | 0.91 | 0.97 | 0.79 | 0.84 | 0.70 | 0.74 | 0.65 | 0.69 |
| SID Meth + Cystine:lysine ^d | Ratio | 0.55 | | 0.55 | | 0.55 | | 0.55 | | 0.55 | |
| SID Threonine:lysine | Ratio | 0.61 | | 0.62 | | 0.63 | | 0.64 | | 0.66 | |
| SID Tryptophan:lysine | Ratio | 0.16 | | 0.16 | | 0.16 | | 0.16 | | 0.16 | |
| SID Valine:lysine | Ratio | 0.68 | | 0.68 | | 0.68 | | 0.68 | | 0.68 | |
| SID Isoleucine:lysine | Ratio | 0.56 | | 0.56 | | 0.56 | | 0.56 | | 0.56 | |
| Calcium | % | 0.75 | | 0.63 | | 0.61 | | 0.58 | | 0.56 | |
| aPhosphorus | % | 0.36 | | 0.30 | | 0.28 | | 0.26 | | 0.22 | |
| Added salt | % | 0.40 | | 0.40 | | 0.40 | | 0.35 | | 0.30 | |
| L-lysine max | % | 0.45 | | 0.40 | | 0.375 | | 0.325 | | 0.225 | |

^a Expected growth and suggested nutrient levels for high health and thermal neutral conditions.

Lysine specifications are based on a series of trials leading to curve validation studies: Technical Memos 160 and 183 and PIC Experiment 9611.

^b Equation used: $0.000027 * \text{weight}^2 - 0.015318 * \text{weight} + 4.114302$

^c Equation used: $(\text{SID lysine per NRC ME} * \text{NRC ME per lb} * 2.2) / 10000$

^d SID = Standardize ileal digestible value.

^e Diet without Paylean®.

Table 19. PIC Barrow Specifications^a

| Barrows | | Phase of growth, lb | | | | | | | | | |
|--|---------|---------------------|------|----------|------|-----------|------|-----------|------|------------------------|------|
| Item | Unit | 50 - 90 | | 90 - 130 | | 130 - 180 | | 180 - 230 | | 230 - 280 ^e | |
| Growth rate | lb/d | 1.82 | | 1.98 | | 2.25 | | 2.15 | | 2.00 | |
| Feed intake | lb/d | 3.15 | | 4.55 | | 5.81 | | 5.88 | | 6.10 | |
| Feed:Gain | Ratio | 1.73 | | 2.30 | | 2.58 | | 2.73 | | 3.05 | |
| NRC ME | Kcal/lb | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 | 1500 | 1600 |
| Added fat | % | 0 | 5 | 0 | 5 | 0 | 5 | 0 | 5 | 0 | 5 |
| SID lysine/Mcal ME ^b | g/Mcal | 3.17 | 3.17 | 2.66 | 2.66 | 2.24 | 2.24 | 1.97 | 1.97 | 1.89 | 1.89 |
| SID lysine ^c | % | 1.05 | 1.12 | 0.88 | 0.94 | 0.74 | 0.79 | 0.65 | 0.69 | 0.62 | 0.67 |
| SID Meth + Cystine:lysine ^d | Ratio | 0.55 | | 0.55 | | 0.55 | | 0.55 | | 0.55 | |
| SID Threonine:lysine | Ratio | 0.61 | | 0.62 | | 0.63 | | 0.64 | | 0.66 | |
| SID Tryptophan:lysine | Ratio | 0.16 | | 0.16 | | 0.16 | | 0.16 | | 0.16 | |
| SID Valine:lysine | Ratio | 0.68 | | 0.68 | | 0.68 | | 0.68 | | 0.68 | |
| SID Isoleucine:lysine | Ratio | 0.56 | | 0.56 | | 0.56 | | 0.56 | | 0.56 | |
| Calcium | % | 0.75 | | 0.63 | | 0.61 | | 0.58 | | 0.56 | |
| aPhosphorus | % | 0.36 | | 0.30 | | 0.28 | | 0.26 | | 0.22 | |
| Added salt | % | 0.40 | | 0.40 | | 0.40 | | 0.35 | | 0.30 | |
| L-lysine max | % | 0.45 | | 0.40 | | 0.375 | | 0.325 | | 0.225 | |

^a Lysine specifications are based on a series of trials leading to curve validation studies:

Technical Memos 160 and 183 and PIC Exp. 9611.

^b Equation used: $0.00004 * \text{weight}^2 - 0.019913 * \text{weight} + 4.369743$

^c Equation used: $(\text{SID lysine per NRC ME} * \text{NRC ME per lb} * 2.2) / 10000$

^d SID = Standardize ileal digestible value.

^e Diet without Paylean®.

Vitamin and Trace Mineral Addition Rates: Commercial^{a,b}

PIC recommendations were established after extensive comparison with Universities and major Nutrition groups. A systematic allowance was made in relation to the NRC recognizing that this information is based on studies under nearly ideal conditions Commercial Nutritionists prior to publication reviewed micronutrient recommendations.

Table 20. Suggested specifications: Amount per pound of complete diet.

| Nutrient | Unit | Nursery | | Grow-Finish | | Sow | Boar |
|--------------------|--------|------------------|------------------|-------------|---------|-------------|------|
| | | < 12 lbs | 12-60 lbs | 60-150 lbs | 150-Mkt | Gest - Lact | Stud |
| Vitamin A | IU/lb | 5000 | 4500 | 3000 | 2200 | 4500 | 5000 |
| Vitamin D | IU/lb | 800 | 750 | 550 | 450 | 800 | 800 |
| Vitamin E | IU/lb | 38 | 35 | 15 | 10 | 30 | 50 |
| Vitamin K | mg/lb | 2.5 | 2.0 | 1.5 | 1.0 | 2.0 | 2.0 |
| Choline | mg/lb | 200 | 150 | 50 | 0 | 300 | 300 |
| Niacin | mg/lb | 32 | 20 | 12 | 10 | 20 | 20 |
| Riboflavin | mg/lb | 6 | 4.5 | 2.6 | 2 | 4.5 | 4.5 |
| d-Pantothenate | mg/lb | 18 | 15 | 9 | 6.5 | 15 | 15 |
| Vitamin B12 | mcg/lb | 25 | 20 | 12 | 10 | 17 | 17 |
| Folic Acid | mcg/lb | 475 | 350 | 0 | 0 | 600 | 750 |
| d-Biotin | mcg/lb | 125 | 70 | 0 | 0 | 100 | 250 |
| Thiamine | mg/lb | 1.6 | 1.5 | 0 | 0 | 1.0 | 1.0 |
| Pyridoxine | mg/lb | 3.2 | 2.0 | 0 | 0 | 1.5 | 1.5 |
| Vitamin C (stable) | mg/lb | 60 | 0 | 0 | 0 | 0 | 60 |
| Zinc | PPM | 150 ^c | 130 ^c | 120 | 70 | 125 | 125 |
| Iron | PPM | 200 ^d | 175 ^d | 80 | 65 | 100 | 100 |
| Manganese | PPM | 50 | 45 | 30 | 25 | 50 | 50 |
| Copper | PPM | 18 ^c | 15 ^c | 12 | 10 | 15 | 15 |
| Iodine | PPM | 0.65 | 0.55 | 0.40 | 0.35 | 0.35 | 0.65 |
| Selenium | PPM | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 |

^a B-Vitamins supplemented at approximately 3.5 x NRC (1998) for < 12 lbs pigs. Multiples for other groups approx. as follows: 12-60 lbs, 3 x NRC.

60-150 lbs, 2.5 x NRC.

150-Mkt, 1.5 x NRC.

Sows tend to be 2.5 x NRC for Vitamins in general.

Boars are similar to sows with extra margins set for several micro-nutrients.

Add 2.3 IU of Vitamin E/lb of complete diet for each 1% fat above 3% total dietary fat.

^b Pelleting and/or expanding decreases Vitamin stability by 10-12% and 15-20% respectively. Consult Vitamin manufacturer to verify the extent by vitamin so additional fortification can be made as required.

^c Nutritional levels are shown for Zinc and Copper. Chemotherapeutic levels of Zinc as follows: < 12 lbs 2600 PPM; 12-16 lbs., 2200 PPM; 16-22 lbs, 1600 PPM. Chemotherapeutic levels of Copper is 220 PPM for each phase. Inorganic forms assumed.

^d Supplemental iron are near to NRC levels because of the substantial iron content of di-calcium phosphate and because high iron intake encourages E.Coli proliferation in the young pig.

Vitamin and Trace Mineral Addition Rates: PIC Multiplication^{a, b}

PIC recommendations were established after extensive comparison with University and major Nutrition groups. A systematic allowance was made relative to the NRC Micronutrient recommendations for Vitamins D and Biotin are contractual for Multiplication. Other micronutrients are recommended minimums. Levels shown assume micronutrient additions and give no credit for ingredient content.

Table 21. Suggested specifications: Amount per pound of complete diet.

| Nutrient | Unit | Nursery | | Grow-Finish | | Sow | Boar |
|--------------------|--------|------------------|------------------|-------------|---------|-------------|------|
| | | < 12 lbs | 12-60 lbs | 60-150 lbs | 150-Mkt | Gest - Lact | Stud |
| Vitamin A | IU/lb | 5000 | 4500 | 3500 | 2600 | 4500 | 5000 |
| Vitamin D | IU/lb | 850 | 800 | 700 | 600 | 800 | 800 |
| Vitamin E | IU/lb | 38 | 35 | 15 | 10 | 30 | 50 |
| Vitamin K | mg/lb | 2.5 | 2.0 | 1.5 | 1.0 | 2.0 | 2 |
| Choline | mg/lb | 200 | 150 | 50 | 0 | 300 | 300 |
| Niacin | mg/lb | 32 | 20 | 12 | 10 | 20 | 20 |
| Riboflavin | mg/lb | 6 | 4.5 | 2.6 | 2 | 4.5 | 4.5 |
| d-Pantothenate | mg/lb | 18 | 15 | 9 | 6.5 | 15 | 15 |
| Vitamin B12 | mcg/lb | 25 | 20 | 12 | 10 | 17 | 17 |
| Folic Acid | mcg/lb | 475 | 350 | 340 | 205 | 600 | 750 |
| d-Biotin | mcg/lb | 150 | 125 | 125 | 90 | 100 | 250 |
| Thiamine | mg/lb | 1.6 | 1.5 | 1.1 | 0.7 | 1.0 | 1.0 |
| Pyridoxine | mg/lb | 3.2 | 2.0 | 1.1 | 0.7 | 1.5 | 1.5 |
| Vitamin C (stable) | mg/lb | 60 | 0 | 0 | 0 | 0 | 60 |
| Zinc | PPM | 135 ^c | 125 ^c | 100 | 70 | 125 | 125 |
| Iron | PPM | 110 ^d | 100 ^d | 80 | 65 | 100 | 100 |
| Manganese | PPM | 40 | 35 | 20 | 16 | 35 | 35 |
| Copper | PPM | 18 ^c | 15 ^c | 12 | 10 | 15 | 15 |
| Iodine | PPM | 0.65 | 0.55 | 0.40 | 0.35 | 0.35 | 0.65 |
| Selenium | PPM | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 |

^a Multiplication VTM requires the addition of greater amounts of Vitamins D and Biotin in Nursery and Grow-Finish diets as compared to commercial specifications. The addition of Folic Acid, Thiamine and Pyridoxine during the Grow-Finish phase is suggested as an extra precaution. Addition rates assume an ME level typical for Corn-Soy diets.

B-Vitamins are supplemented at about 3.5 x NRC (1998) for < 12 lbs pigs. Multiples for other groups approx. as follows: 12-60 lbs, 3 x NRC.

60-150 lbs, 2.5 x NRC.

150-Mkt, 1.5 x NRC.

Sows tend to be 2.5 x NRC for Vitamins in general.

Boars are similar to sows with extra margins set for several micronutrients.

Add 2.3 IU of Vitamin E / lb of complete diet for each 1% fat above 3% total dietary fat.

^b Pelleting and/or expanding decreases Vitamin stability by 10-12% and 15-20% respectively. Consult Vitamin manufacturer to verify the extent by vitamin so additional fortification can be made as required.

^c Nutritional levels are shown for Zinc and Copper. Chemotherapeutic levels of Zinc as follows: < 12 lbs 2600 PPM; 12-16 lbs., 2200 PPM; 16-22 lbs, 1600 PPM. Chemotherapeutic level for Copper is 225 PPM for each phase. Inorganic forms assumed. Sulfate forms of Zn, Fe, Mn and Cu are preferred.

^d Supplemental iron are near to NRC levels because of the substantial iron content of di-calcium phosphate and because high iron intake encourages E.Coli proliferation in the young pig.

Alternative ingredients

With the price increase in corn and soybean meal many producers are looking at using alternative ingredients like corn dried distiller's grains, bakery by-products, glycerin and fractioned corn. Please remember that these products are by-products and users must be careful when using them. Some alternative ingredients are not well studied and should be used with extreme caution. Glycerin is still being studied for an energy source and therefore the use should be limited with a max of 6% because of knowledge and feed flowability. These products tend to be very variable in nutrients and may contain mycotoxins. Ingredient samples must be taken and analyzed to determine their nutrient levels. Consistency is very important so try and work with one plant when receiving alternative ingredients to get a consistent product. When dealing with alternative ingredients producers must keep in mind feed mill space, feed flowability, and carcass traits.

There are two feeding strategies for DDGS they are 1) Step up program, 2) High inclusion. The idea of a step up program is to slowly let the pigs get used to the taste and smell of DDGS and not adding a lot of DDGS suddenly. The second idea is to have a high level of DDGS in the starter diets so the pigs will have to get used to DDGS early and use more DDGS faster. New research from Gaines et al. (2007), indicates decreased carcass yield with high levels of DDGS due to increased intestine weight. Other ingredients that may decrease yield if used at a too high of level are soybean meal, wheat midds and soy hulls.

Some packers are now testing for iodine value (IV) which is a measure of the level of unsaturation or softness of a fat. Benz et al. (2007a,c) has proven that IV will increase with inclusion of oils with high levels of unsaturated fatty acids like corn oil from DDGS. Other ingredients that will increase IV are extruded expelled soybean meal (Benz et al., 2007b) and high oil corn. Below is an example of the amount (%) of DDGS that can be added to nursery and finishing diets that will minimize the yield decline and keep the IV low (Table 22). This guide also shows when new ingredients are used that only 5% increases should be done. Stepping up faster with DDGS or other by-products can result in lower feed intakes. Other ingredients in the diet can also affect IV so consultant with your nutritionist and packer on IV specs. Breeding herd inclusions are shown in Table 23.

Research also has proved that when dietary linoleic acid is increased this will cause IV to increase.

Producers must work closely with their nutritionists to implement these ingredients in their diets.


Table 22. Amount of DDGS in diets for market pigs or replacement gilts.

| DDGS | Unit | 12 to 15 | 15 to 25 | 25 to 50 | 50 to 90 | 90 to 150 | 150 to 210 | 210 to 280 | Paylean |
|----------------|------|----------|----------|----------|----------|-----------|------------|------------|---------|
| Market | % | 0.0 | 5.0 | 10.0 | 15.0 | 15.0 | 10.0 | 0.0 | 0.0 |
| Multiplication | % | 0.0 | 5.0 | 10.0 | 15.0 | 15.0 | 15.0 | 15.0 | -- |

Table 23. Amount of DDGS in the breeding herd.

| Diet | Unit | DDGS |
|-----------|------|---------|
| Gestation | % | 0 to 30 |
| Lactation | % | 0 to 20 |
| Boar | % | 0 to 15 |

Potential issues with feeding DDGS



Pork Quality

Below are pictures courtesy of Purdue University from bellies from pigs that were fed either 0%, 10%, 20%, or 30% DDGS. The pictures show the flop test and as higher levels of DDGS was fed the fat on the bellies got softer.

0% DDGS



10% DDGS



20% DDGS



30% DDGS



Paylean®^a

Paylean® is a great product with proven results when diets are formulated correctly. With the new labeling in 2006 with Paylean®, market pigs can be fed to heavier weights. Please keep in mind that Paylean® should not be fed to replacement gilts or boars. Below is a guideline when feeding Paylean® for less than 21 days or greater than 21. With the SID lysine in the Paylean diet being high there is a risk of adding too much soybean meal and causing a yield reduction (Gaines et al., 2004 and 2007).

Table 24. Diet specifications for diets with Paylean®

| Paylean® | Unit | < 21 days | > 21 days |
|---------------------------|---------|-----------|-----------|
| Growth rate | lb/day | 2.25 | 2.15 |
| Feed intake | lb/day | 6.10 | 6.20 |
| Feed:Gain ratio | F/G | 2.71 | 2.88 |
| NRC ME | kcal/lb | 1550 | 1550 |
| SID Lysine | % | 0.95 | 0.85 |
| Added Fat | % | 2 to 3 | 2 to 3 |
| SID Meth + Cystine:lysine | Ratio | 0.55 | |
| SID Threonine:lysine | Ratio | 0.68 | |
| SID Tryptophan:lysine | Ratio | 0.16 | |
| SID Valine:lysine | Ratio | 0.68 | |
| SID Isoleucine:lysine | Ratio | 0.56 | |
| Calcium | % | 0.56 | |
| T Phosphorus | % | 0.51 | |
| aPhosphorus | % | 0.24 | |
| Added salt | % | 0.30 | |
| Min L-lysine | % | 0.30 | |
| Max L-lysine | % | 0.40 | |

^a Paylean® is Elanco's trade mark for ractopamine HCl.

Effects of nutrition on yield

Recent research has demonstrated that ingredient type can affect the yield of PIC finishing pigs. As mentioned in the alternative ingredient section, high fiber ingredients have shown to decrease yield. Also in the Paylean® section references low soybean meal and high synthetic amino acid diets can increase yield. High levels of soybean meal throughout the finishing period can also cause low yield. Below is a trial conducted with high levels of soybean meal and low levels of synthetic amino acids (Tables 25 and 26).

Table 25. Effects of soybean meal on carcass yield^a

| Diets | SBM levels | | | P - value |
|--------------------|------------|--------|-------|-----------|
| | Low | Medium | High | Linear |
| Corn, % | 69.8 | 64.5 | 59.2 | -- |
| SBM, % | 22.5 | 27.6 | 32.7 | -- |
| L-Lysine, % | 0.225 | 0.225 | 0.225 | -- |
| Results | | | | |
| Carcass weight, lb | 222.4 | 218.4 | 215.9 | 0.05 |
| Carcass yield, % | 76.5 | 76.2 | 74.4 | 0.09 |

^a Pigs were PIC 337 x Camborough with 8 replications per treatment.

Table 26. Effect of low versus high synthetic amino acid diets for PIC gilts fed Paylean® (6.5 g/ton) for 21 days^a

| Item | L-Lysine, lbs/ton | | SEM | P-Value |
|------------------|------------------------------|----------------------------|------|---------|
| | 1.5 lbs (556 lbs/ton SBM) | 6 lbs (398 lbs/ton SBM) | | |
| Growth | | | | |
| Day 0 BW, lbs | 229.7 | 230.8 | 1.30 | 0.57 |
| Day 21 BW, lbs | 278.0 | 278.1 | 1.50 | 0.96 |
| ADG, lbs/d | 2.30 | 2.25 | 0.03 | 0.33 |
| ADFI, lbs/d | 6.20 | 6.08 | 0.08 | 0.32 |
| Feed/Gain | 2.70 | 2.70 | 0.04 | 0.96 |
| Carcass | | | | |
| Carcass wt., lbs | 207.1 | 211.4 | 1.07 | 0.03 |
| Backfat, in. | 0.64 | 0.66 | 0.02 | 0.63 |
| Yield, % | 76.4 | 77.6 | 0.38 | 0.05 |
| Lean, % | 56.0 | 55.8 | 0.29 | 0.69 |

^aData represents the means of 8 replicate pens (21 pigs/pen). Diets were formulated to a 0.95% SID lysine (2.78 g SID lysine/Mcal ME) and fed for 21 days. Trial was conducted at PorkTech, LLC (Moberly, MO).

Feed Manufacturing

Guidelines for feed manufacturing for PIC pigs.

Meal feed:

Roller mill

700 microns with a standard deviation of < 2.3

Mixing CV of less than 10%

- < 30 % over 1 mm
- < 20 % under 300 microns
- < 2.5 % under 150 microns

Pellets:

Hammer mill

500 microns with a standard deviation of < 2.5

PDI > 90

Feed Form:

With increased feed prices feed efficiency has become increasingly important. PIC has done a number of large scale trials comparing feed form of meal and pellets on feed efficiency. Below are a few trials comparing sire lines and meal versus pellets on growth performance (Table 27, 28 and 29). The trial results demonstrate that feeding pellets will improve (6% to 10%) feed efficiency in all sire lines however full value pigs may be higher (1% to 3%) in meal diets for some lines. All pigs were not vaccinated for PCV2 and not fed Paylean®.

Table 27. Impact of feed form on PIC 280 sired pigs^a

| Item | Sire line x Feed form | | P < |
|---------------------|-----------------------|----------------|--------|
| | PIC 280 Meal | PIC 280 Pellet | |
| On test weight, lb | 62.9 | 64.3 | 0.53 |
| Off test weight, lb | 255.1 | 264.8 | 0.016 |
| ADG, lb | 2.03 | 2.12 | 0.0011 |
| F/G | 2.74 | 2.48 | 0.0001 |
| Age at 275 lb, d | 174.4 | 172.1 | 0.0001 |
| Survival rate, % | 97.2 | 98.2 | 0.27 |
| Yield, % | 74.5 | 74.6 | 0.19 |
| Lean, % | 54.7 | 54.5 | 0.23 |

^a A total of 1,044 pigs (PIC 280 x Camborough) were used with 41 replications for meal and 43 replications for pellets.

Table 28. Impact of feed form on PIC 337 sired pigs^a

| Item | Sire line x Feed form | | P < |
|---------------------|-----------------------|----------------|--------|
| | PIC 337 Meal | PIC 337 Pellet | |
| On test weight, lb | 53.3 | 53.2 | 0.97 |
| Off test weight, lb | 265.3 | 269.3 | 0.01 |
| ADG, lb | 2.08 | 2.12 | 0.01 |
| F/G | 2.52 | 2.36 | 0.0001 |
| Age at 275 lb, d | 171.0 | 168.9 | 0.0003 |
| Deads, % | 2.1 | 3.1 | 0.18 |
| Culls, % | 3.5 | 4.1 | 0.52 |
| Lean, % | 56.1 | 55.6 | 0.0004 |

^a A total of 1,730 pigs (PIC 337 x Camborough) were used with 75 replications for meal and 73 replications for pellets.

Table 29. Impact of feed form on PIC 380 sired pigs^a

| Item | Feed form | | P < |
|---------------------|-----------|--------|--------|
| | Meal | Pellet | |
| On test weight, lb | 55.7 | 54.5 | 0.30 |
| Off test weight, lb | 268.2 | 270.3 | 0.22 |
| ADG, lb | 2.11 | 2.15 | 0.003 |
| F/G | 2.49 | 2.36 | 0.0001 |
| Age at 275 lb, d | 168.8 | 166.8 | 0.0001 |
| Deads, % | 1.5 | 3.4 | 0.01 |
| Culls, % | 1.5 | 2.9 | 0.06 |
| Backfat, in | 0.68 | 0.71 | 0.0001 |
| Loin depth, in | 2.31 | 2.36 | 0.0001 |
| Lean, % | 55.4 | 55.1 | 0.009 |

^a A total of 1,699 pigs (PIC 380 x Camborough) were used with 76 replications for meal and 74 replications for pellets.

Feeder Space

Guidelines for feeder space for PIC pigs depending on feeder type.
Feeder Requirements:

Wet or Dry Feeders:

- Maximum of 12 pigs per feeder space
- Minimum of 14 inches per head space

Tube Feeders:

- Maximum of 10 pigs per tube
- Minimum of 2 inches trough space per pig

References

- Benz, J.M., M.D. Tokach, S.S. Dritz, J.L. Nelssen, J.M. DeRouche, and R.D. Goodband. 2007a. Effects of choice white grease or soybean oil on growth performance and carcass characteristics of grow-finish pigs. *Kansas Agric. Exp. Stn. Rep. Prog. No. 985*. pp. 99-104.
- Benz, J.M., M.D. Tokach, S.S. Dritz, J.L. Nelssen, J.M. DeRouche, and R.D. Goodband. 2007b. Effects of DDGS and extruded expelled soybean meal on growth performance and carcass characteristics of grow-finish pigs. *Kansas Agric. Exp. Stn. Rep. Prog. No. 985*. pp. 114-121.
- Benz, J.M., S.K. Linneen, J.M. DeRouche, M.D. Tokach, S.S. Dritz, J.L. Nelssen, and R.D. Goodband. 2007c. Effects of dried distillers grains with solubles on growth performance and fat quality of finishing pigs. *Kansas Agric. Exp. Stn. Rep. Prog. No. 985*. pp. 105-113.
- Boyd, R.D., K.J. Touchette, G.C. Castro, M.E. Johnston, K.U. Lee and In K. Han. 2000a. Recent Advances in amino acid and energy nutrition of prolific sows. *Asian-Aus. J. Anim. Sci.* Vol. 13, No. 11:1638-1652.
- Boyd, R.D., M.E. Johnston, and G. Castro. 2000b. Feeding to achieve genetic potential. Banff Conference.
- Boyd, R.D., M.E. Johnston, J.L. Usry, C.E. Fralick, A.A. Sosnicki and B. Fields. 2001. Lysine level required to optimize the growth performance to Paylean in PIC pigs. *J. Anim. Sci.* 79 (Suppl. 1):66 (Abstr.).
- Boyd, R.D., G.C. Castro, R.A. Cabrera. 2002. Nutritional management of the sow to maximize lifetime productivity. Banff Conference.
- Boyd, R.D., N.H. Williams. 2008. Segregated parity structure in sow farms to capture nutrition, management and health opportunities. Manitoba Swine Seminar.
- Gaines, A.M., B.W. Ratiff, P. Srichana, G.L. Allee and J.L. Usry. 2004. Evaluation of high synthetic lysine diets for pigs fed ractopamine HCl (Paylean). *J. Anim. Sci.* 82 (Suppl.2):38 (Abstr.).
- Gaines, A.M. J.D. Spencer, G.I. Petersen, N.R. Augspurger and S.K. Kitt. 2007. Effect of corn distiller's dried grains with solubles (DDGS) withdrawal program on growth performance and carcass yield in grow-finish pigs.
- Gaines, A.M., R.D. Boyd, M.E. Johnston, G.L. Allee, and J.L. Usry. 2007. Lysine source affects ractopamine diets. *Feedstuffs*. Vol. 79, No. 16, March 26, 2007.
- Hastad, C.W., S.S. Dritz, M.D. Tokach, R.D. Goodband, J.L. Nelssen, J.M. DeRouche, R.D. Boyd, M.E. Johnston. 2004. Phosphorus requirements of growing-finishing pigs reared in a commercial environment. *J. Anim. Sci.* 82:2945-2952.
- Neill, C. R., S.S. Dritz, M.D. Tokach, J.L. Nelssen, R.D. Goodband, J.M. DeRouche and J.L. Usry. 2006. Lysine requirement of pigs fed ractopamine HCl in a commercial facility. *J. Anim. Sci.* 84 (Suppl. 2):197 (Abstr.)
- Nutrient Requirements of Swine. 1998. National Research Council (10th Edition). National Academy Press. 2101 Constitution Ave., NW. Washington, D.C. 20418.
- Optimizing energy formulation for finishing swine. 1997. J. Usry, R.G. Campbell and D. Burnham. Proc. Carolina Swine Nutrition Conference, Raleigh NC.
- Performance of 25 to 60 lb pigs is optimized at 58% TID M+C:Lysine. Ajinomoto Heartland LLC Swine Research Report 38.
- PIC Fundamentals of gilt and sow management. 2007.
- PIC Technical Memo 142. 1996. Impact of dietary energy intake on sperm output by PIC 327 and PIC 427 boars.
- PIC Technical Memo 160. 1997. Comparison of two dietary lysine standards for growing PIC pigs: PIC vs University of Illinois recommendations.
- PIC Technical Memo 166. 1997. Response to three lysine regimens by PIC 406 progeny.
- PIC Technical Memo 171. 1997. Amino acid requirements of the lactating sow: Literature Review. Adapted and improved from K. Touchette MS Thesis, University of Missouri.

- PIC Technical Memo 183. 1997. Dietary lysine curve that optimizes lean growth and profit for PIC 427 progeny.
- PIC Technical Memo 153. 1997. Relationship between dietary fatty acid profile and body fat composition in growing pigs.
- PIC Experiment 9611. Lysine dose response curves for heavy PIC pigs (210-260 lbs) reared in a hot vs thermo-neutral environment.
- PIC Nutrition Technical Update. 1999. Concepts for feeding the Grow-Finish PIC pig.
- PIC Nutrition Technical Update. 1999. Concepts for feeding prolific PIC sows.
- PIC Technical Memo 245. 2001. Response of PIC 337 progeny to the metabolism modifier Paylean®: Response is dependent on dietary lysine intake.
- PIC Technical Memo 10. 2004. Effects of space allowance and feed type on grow – finish performance of PIC 280 sired pigs.
- PIC Technical Memo 330. 2007. Grow – Finish Performance of PIC 380 Sired pigs: How is it effected by space allowance and feed form.
- PIC Technical Memo 331. 2007. Grow – Finish Performance of PIC 337 sired pigs: Effects of space allowance and feed form.
- Ratliff, B., G. Allee, A. Gaines, B. Hinson. 2007. Maximizing the use of feed grade amino acids. Ajinomoto Heartland LLC and PIC Swine Summit.
- Srichana, Pairat. 2006. Amino Acid Nutrition in Gestation and Lactating Sows. PhD Thesis. University of Missouri.
- Threonine:Lysine ratio for optimal performance of late finishing swine. Ajinomoto Heartland LLC Swine Research Report 36.
- Williams, N, R. Kummer, J. Pinilla, J. Piva, C. Neill. 2007. Milk production and nutritional requirements in modern sows. ABRAVES. Association of Brazilian Veterinarians.

A special thank you goes to Dr. Dean Boyd, Dr. James Usry, and Dr. Gary Allee for their help in data and research results along with review of this nutrition guide.

DISCLAIMER. Performance data shown in Nutrient specification tables were obtained in Commercial settings and under conditions of high health, thermo-neutral temperature and good management. They are not guaranteed levels of performance. A competent Nutritionist should adapt suggested Nutrient levels to specific conditions. These concepts are discussed in greater detail in Nutrition Technical Updates for Sows, Nursery pigs and Grow-Finish pigs.

PIC
100 Bluegrass Commons Blvd.
Ste. 2200
Hendersonville, TN 37075
www.pic.com/usa
1-800-325-3398

Copyright 2008 PIC All rights reserved.

